

WEST Search History

DATE: Thursday, November 06, 2003

Set Name Query

side by side

Hit Count Set Name

result s t

DB=USPT; PLUR=YES; OP=AND

L1	(helicobacter or helicoacter or hpyhori or hpylori or pylori or pyloris or pyloridis).clm.	410	L1
L2	L1 and urease.clm.	54	L2
L3	L2 and (weight or kda or kd or da or daltons or dalton or rmw or size).clm.	11	L3
L4	(hsp or hspb or hsp-b).clm.	79	L4
L5	L4 and l1	4	L5

END OF SEARCH HISTORY

WEST Search History

DATE: Thursday, November 06, 2003


<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
side by side			result set

DB=USPT; PLUR=YES; OP=AND

L1	hspa.clm. and hspb.clm.	3	L1
----	-------------------------	---	----

L2	hspa.clm. and ureb.clm.	2	L2
----	-------------------------	---	----

END OF SEARCH HISTORY

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot			
Hosted by NCSC US	Mirror sites:	Bolivia	Canada	China	Korea	Switzerland	Taiwan

Search for 

11/03/03

Search in Swiss-Prot and TrEMBL for: helicobacter pylori kda

Swiss-Prot Release 42.1 of 24-Oct-2003

TrEMBL Release 25.1 of 24-Oct-2003

Number of sequences found in Swiss-Prot₍₁₇₎ and TrEMBL₍₁₇₎: **34**

Note that the selected sequences can be saved to a file to be later retrieved; to do so, go to the bottom of this page.

For more directed searches, you can use the Sequence Retrieval System SRS.

Search in Swiss-Prot: There are matches to 17 out of 136356 entries

AN36_HELPJ (Q9ZJD1)

36 kDa antigen. {GENE: JHP1381} - Helicobacter pylori J99
(Campylobacter pylori J99)

AN36_HELPY (P94851)

36 kDa antigen. {GENE: HP1488} - Helicobacter pylori (Campylobacter

pylori)

CAGA_HELPJ (Q9ZLT1)

Cytotoxicity associated immunodominant antigen (120 kDa protein) (CAG pathogenicity island protein 26). {GENE: CAGA OR CAI OR CAG26 OR JHP0495} - Helicobacter pylori J99 (Campylobacter pylori J99)

CAGA_HELPY (P55980)

Cytotoxicity associated immunodominant antigen (120 kDa protein) (CAG pathogenicity island protein 26). {GENE: CAGA OR CAI OR CAG26 OR HP0547} - Helicobacter pylori (Campylobacter pylori)

CGA1_HELPY (P80200)

Cytotoxicity associated immunodominant antigen (120 kDa protein). {GENE: CAGA OR CAI} - Helicobacter pylori (Campylobacter pylori)

CGA2_HELPY (P55746)

Cytotoxicity associated immunodominant antigen (120 kDa protein). {GENE: CAGA OR CAI} - Helicobacter pylori (Campylobacter pylori)

CH10_HELPY (P48225)

10 kDa chaperonin (Protein Cpn10) (groES protein) (Heat shock protein 10). {GENE: GROS OR GROES OR MOPB OR HSP10 OR HSPA OR HP0011 OR JHP0009} - Helicobacter pylori (Campylobacter pylori), Helicobacter pylori J99 (Campylobacter pylori J99)

CH60_HELPJ (Q9ZN50)

60 kDa chaperonin (Protein Cpn60) (groEL protein) (Heat shock protein 60). {GENE: GROL OR MOPA OR GROEL OR HSP60 OR HSPB OR JHP0008} - Helicobacter pylori J99 (Campylobacter pylori J99)

CH60_HELPY (P42383)

60 kDa chaperonin (Protein Cpn60) (groEL protein) (Heat shock protein 60). {GENE: GROL OR MOPA OR GROEL OR HSP60 OR HSPB OR HP0010} - Helicobacter pylori (Campylobacter pylori)

DNAK_HELPJ (Q9ZMW4)

Chaperone protein dnaK (Heat shock protein 70) (Heat shock 70 kDa protein) (HSP70). {GENE: DNAK OR JHP0101} - Helicobacter pylori J99 (Campylobacter pylori J99)

DNAK_HELPY (P55994)

Chaperone protein dnaK (Heat shock protein 70) (Heat shock 70 kDa protein) (HSP70). {GENE: DNAK OR HP0109} - Helicobacter pylori

(Campylobacter pylori)

HCPA_HELPJ (Q9ZMM1)

Beta-lactamase hcpA precursor (EC 3.5.2.6) (Cysteine-rich 28 kDa protein).
{GENE: HCPA OR JHP0197} - Helicobacter pylori J99 (Campylobacter pylori J99)

HCPA_HELPY (O25001)

Beta-lactamase hcpA precursor (EC 3.5.2.6) (Cysteine-rich 28 kDa protein).
{GENE: HCPA OR HP0211} - Helicobacter pylori (Campylobacter pylori)

PYRX_HELPJ (Q9ZMG9)

Probable dihydroorotase-like protein (Aspartate carbamoyltransferase 42 kDa non-catalytic chain). {GENE: PYRC' OR JHP0251} - Helicobacter pylori J99 (Campylobacter pylori J99)

PYRX_HELPY (O25045)

Probable dihydroorotase-like protein (Aspartate carbamoyltransferase 42 kDa non-catalytic chain). {GENE: PYRC' OR HP0266} - Helicobacter pylori (Campylobacter pylori)

TSAA_HELPJ (P56876)

Probable peroxiredoxin (26 kDa antigen). {GENE: TSAA OR JHP1471} - Helicobacter pylori J99 (Campylobacter pylori J99)

TSAA_HELPY (P21762)

Probable peroxiredoxin (26 kDa antigen). {GENE: TSAA OR HP1563} - Helicobacter pylori (Campylobacter pylori)

Search in TrEMBL: There are matches to 17 out of 1016356 entries

O25989

60 kDa inner-membrane protein {GENE:HP1450} - Helicobacter pylori (Campylobacter pylori)

O50170

Putative 32.7 kDa protein - Helicobacter pylori (Campylobacter pylori)

P94847

48 kDa antigen - Helicobacter pylori (Campylobacter pylori)

P94848

29 kDa protein - Helicobacter pylori (Campylobacter pylori)

Q8RNU2

Heat shock protein B subunit (60 kDa chaperonin) (Protein Cpn60) (groEL protein) {GENE:HSPB} - Helicobacter pylori (Campylobacter pylori)

Q8RNU3

Heat shock protein A subunit (10 kDa chaperonin) (Protein Cpn10) (groES protein) {GENE:HSPA} - Helicobacter pylori (Campylobacter pylori)

Q8RNU7

20 kDa membrane-associated lipoprotein (Membrane-associated lipoprotein Lpp20) {GENE:LPP20} - Helicobacter pylori (Campylobacter pylori)

Q93SI8

26,000 kDa protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R2X6

60 kDa major heat shock protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R4J2

31 kDa major NONSELECTIVE porin protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R4W4

25 kDa major heat shock protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R4W5

30 kDa major heat shock protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R4W6

56 kDa major heat shock protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R4W7

66 kDa major heat shock protein (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R510

60 kDa heat shock protein/groEL homolog (Fragment) - Helicobacter pylori (Campylobacter pylori)

Q9R5L5

HSP62=62 kDa urease-associated heat shock protein (Fragment) -
Helicobacter pylori (Campylobacter pylori)

Q9X749

HopV protein precursor (27 kDa outer membrane protein) {GENE:HOPV OR
OMP27} - Helicobacter pylori (Campylobacter pylori)

New Search


in Swiss-Prot/TrEMBL by AC, ID,
description, gene name, organism
Please do NOT use any boolean
perators (and, or, etc.)


If you would like to retrieve all the Swiss-Prot/TrEMBL entries contained in
this list , you can enter a file name. These entries will then be saved to a file
under this name in the directory outgoing of the ExpASy anonymous ftp server,
from where you can download it. (Please note that this temporary file will only
be kept for 1 week.)

File name:

Format: ☒ Swiss-Prot ☐ Fasta

or

 ExpASy Home page	Site Map	Search ExpASy	Contact us	Swiss-Prot			
Hosted by NCSC US	Mirror sites:	Bolivia	Canada	China	Korea	Switzerland	Taiwan

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot			
Hosted by NCSC US	Mirror sites:	Bolivia	Canada	China	Korea	Switzerland	Taiwan
Search		<input type="text" value="Swiss-Prot/TrEMBL"/>	<input type="button" value="Go"/>	<input type="button" value="Clear"/>			

Search in Swiss-Prot and TrEMBL for: helicobacter pylori da

Swiss-Prot Release 42.1 of 24-Oct-2003
TrEMBL Release 25.1 of 24-Oct-2003

Number of sequences found in Swiss-Prot₍₈₅₎ and TrEMBL₍₉₅₎: **180**

Note that the selected sequences can be saved to a file to be later retrieved; to do so, go to the bottom of this page.

For more directed searches, you can use the Sequence Retrieval System SRS.

Search in Swiss-Prot: There are matches to 85 out of 136356 entries

AMIE_HELPJ (Q9ZME1)

Aliphatic amidase (EC 3.5.1.4) (Acylamide amidohydrolase). {GENE: AMIE *37 kDa*
OR JHP0279} - Helicobacter pylori J99 (Campylobacter pylori J99)

AMIE_HELPY (O25067)

Aliphatic amidase (EC 3.5.1.4) (Acylamide amidohydrolase). {GENE: AMIE *37 kDa*

OR HP0294} - Helicobacter pylori (Campylobacter pylori)

AMIF_HELPJ (Q9ZJY8)

Formamidase (EC 3.5.1.49) (Formamide amidohydrolase). {GENE: AMIF OR JHP1159} - Helicobacter pylori J99 (Campylobacter pylori J99) 37kDa

AMIF_HELPY (O25836)

Formamidase (EC 3.5.1.49) (Formamide amidohydrolase). {GENE: AMIF OR HP1238} - Helicobacter pylori (Campylobacter pylori) 37kDa

AMPA_HELPJ (Q9ZLR1)

Cytosol aminopeptidase (EC 3.4.11.1) (Leucine aminopeptidase) (LAP) (Leucyl aminopeptidase). {GENE: PEPA OR JHP0517} - Helicobacter pylori J99 (Campylobacter pylori J99)

AMPA_HELPY (O25294)

Cytosol aminopeptidase (EC 3.4.11.1) (Leucine aminopeptidase) (LAP) (Leucyl aminopeptidase). {GENE: PEPA OR HP0570} - Helicobacter pylori (Campylobacter pylori)

AMPM_HELPJ (Q9ZJT0)

Methionine aminopeptidase (EC 3.4.11.18) (MAP) (Peptidase M). {GENE: MAP OR JHP1219} - Helicobacter pylori J99 (Campylobacter pylori J99)

AMPM_HELPY (P56102)

Methionine aminopeptidase (EC 3.4.11.18) (MAP) (Peptidase M). {GENE: MAP OR HP1299} - Helicobacter pylori (Campylobacter pylori)

AN36_HELPJ (Q9ZJD1)

36 kDa antigen. {GENE: JHP1381} - Helicobacter pylori J99 (Campylobacter pylori J99) 36kDa

AN36_HELPY (P94851)

36 kDa antigen. {GENE: HP1488} - Helicobacter pylori (Campylobacter pylori) 36kDa

BIOA_HELPJ (Q9ZKM5)

Adenosylmethionine-8-amino-7-oxononanoate aminotransferase (EC 2.6.1.62) (7,8-diamino-pelargonic acid aminotransferase) (DAPA aminotransferase). {GENE: BIOA OR JHP0910} - Helicobacter pylori J99 (Campylobacter pylori J99) 49,944 Da

BIOA_HELPY (O25627)

Adenosylmethionine-8-amino-7-oxononanoate aminotransferase (EC 2.6.1.62) (7,8-diamino-pelargonic acid aminotransferase) (DAPA 49,768 Da

aminotransferase). {GENE: BIOA OR HP0976} - Helicobacter pylori
(Campylobacter pylori)

CAGA_HELPJ (Q9ZLT1)

Cytotoxicity associated immunodominant antigen (120 kDa protein) (CAG pathogenicity island protein 26). {GENE: CAGA OR CAI OR CAG26 OR JHP0495} - Helicobacter pylori J99 (Campylobacter pylori J99)

CAGA_HELPY (P55980)

Cytotoxicity associated immunodominant antigen (120 kDa protein) (CAG pathogenicity island protein 26). {GENE: CAGA OR CAI OR CAG26 OR HP0547} - Helicobacter pylori (Campylobacter pylori)

CDSA_HELPJ (Q9ZML7)

Phosphatidate cytidylyltransferase (EC 2.7.7.41) (CDP-diglyceride synthetase) (CDP-diglyceride pyrophosphorylase) (CDP-diacylglycerol synthase) (CDS) (CTP:phosphatidate cytidylyltransferase) (CDP-DAG synthase) (CDP-DG synthetase). {GENE: CDSA OR JHP0201} - Helicobacter pylori J99 (Campylobacter pylori J99)

CDSA_HELPY (O25004)

Phosphatidate cytidylyltransferase (EC 2.7.7.41) (CDP-diglyceride synthetase) (CDP-diglyceride pyrophosphorylase) (CDP-diacylglycerol synthase) (CDS) (CTP:phosphatidate cytidylyltransferase) (CDP-DAG synthase) (CDP-DG synthetase). {GENE: CDSA OR HP0215} - Helicobacter pylori (Campylobacter pylori)

CGA1_HELPY (P80200)

Cytotoxicity associated immunodominant antigen (120 kDa protein). {GENE: CAGA OR CAI} - Helicobacter pylori (Campylobacter pylori)

CGA2_HELPY (P55746)

Cytotoxicity associated immunodominant antigen (120 kDa protein). {GENE: CAGA OR CAI} - Helicobacter pylori (Campylobacter pylori)

CH10_HELPY (P48225)

10 kDa chaperonin (Protein Cpn10) (groES protein) (Heat shock protein 10). {GENE: GROS OR GROES OR MOPB OR HSP10 OR HSPA OR HP0011 OR JHP0009} - Helicobacter pylori (Campylobacter pylori), Helicobacter pylori J99 (Campylobacter pylori J99)

CH60_HELPJ (Q9ZN50)

60 kDa chaperonin (Protein Cpn60) (groEL protein) (Heat shock protein

60). {GENE: GROL OR MOPA OR GROEL OR HSP60 OR HSPB OR JHP0008} - Helicobacter pylori J99 (Campylobacter pylori J99)

CH60_HELPY (P42383)

60 kDa chaperonin (Protein Cpn60) (groEL protein) (Heat shock protein

60). {GENE: GROL OR MOPA OR GROEL OR HSP60 OR HSPB OR HP0010} - Helicobacter pylori (Campylobacter pylori)

CLPP_HELPJ (Q9ZL50)

ATP-dependent Clp protease proteolytic subunit (EC 3.4.21.92)

(Endopeptidase Clp). {GENE: CLPP OR JHP0730} - Helicobacter pylori J99 (Campylobacter pylori J99)

CLPP_HELPY (P56156)

ATP-dependent Clp protease proteolytic subunit (EC 3.4.21.92)

(Endopeptidase Clp). {GENE: CLPP OR HP0794} - Helicobacter pylori (Campylobacter pylori)

CLPS_HELPJ (Q9ZN32)

ATP-dependent Clp protease adaptor protein clpS. {GENE: CLPS OR JHP0028} - Helicobacter pylori J99 (Campylobacter pylori J99)

CLPS_HELPY (P56066)

ATP-dependent Clp protease adaptor protein clpS. {GENE: CLPS OR HP0032} - Helicobacter pylori (Campylobacter pylori)

DAPA_HELPJ (Q9ZM13)

Dihydrodipicolinate synthase (EC 4.2.1.52) (DHDPs). {GENE: DAPA OR JHP0410} - Helicobacter pylori J99 (Campylobacter pylori J99)

DAPA_HELPY (O25657)

Dihydrodipicolinate synthase (EC 4.2.1.52) (DHDPs). {GENE: DAPA OR HP1013} - Helicobacter pylori (Campylobacter pylori)

DAPB_HELPJ (Q9ZLW6)

Dihydrodipicolinate reductase (EC 1.3.1.26) (DHPR). {GENE: DAPB OR JHP0460} - Helicobacter pylori J99 (Campylobacter pylori J99)

DAPB_HELPY (P94844)

Dihydrodipicolinate reductase (EC 1.3.1.26) (DHPR). {GENE: DAPB OR HP0510} - Helicobacter pylori (Campylobacter pylori)

DAPF_HELPJ (Q9ZLR5)

Diaminopimelate epimerase (EC 5.1.1.7) (DAP epimerase). {GENE: DAPF OR JHP0513} - Helicobacter pylori J99 (Campylobacter pylori J99)

DAPF_HELPY (O25290)

Diaminopimelate epimerase (EC 5.1.1.7) (DAP epimerase). {GENE: DAPF OR HP0566} - Helicobacter pylori (Campylobacter pylori)

DCDA_HELPJ (Q9ZME5)

Diaminopimelate decarboxylase (EC 4.1.1.20) (DAP decarboxylase). {GENE: LYSA OR JHP0275} - Helicobacter pylori J99 (Campylobacter pylori J99)

DCDA_HELPY (P56129)

Diaminopimelate decarboxylase (EC 4.1.1.20) (DAP decarboxylase). {GENE: LYSA OR HP0290} - Helicobacter pylori (Campylobacter pylori)

DNAK_HELPJ (Q9ZMW4)

Chaperone protein dnaK (Heat shock protein 70) (Heat shock 70 kDa protein) (HSP70). {GENE: DNAK OR JHP0101} - Helicobacter pylori J99 (Campylobacter pylori J99)

DNAK_HELPY (P55994)

Chaperone protein dnaK (Heat shock protein 70) (Heat shock 70 kDa protein) (HSP70). {GENE: DNAK OR HP0109} - Helicobacter pylori (Campylobacter pylori) *Kds*

FLAV_HELPJ (Q9ZK53)

Flavodoxin. {GENE: FLDA OR JHP1088} - Helicobacter pylori J99 (Campylobacter pylori J99)

FLAV_HELPY (O25776)

Flavodoxin. {GENE: FLDA OR HP1161} - Helicobacter pylori (Campylobacter pylori)

FRDA_HELPJ (Q9ZMP0)

Fumarate reductase flavoprotein subunit (EC 1.3.99.1). {GENE: FRDA OR JHP0178} - Helicobacter pylori J99 (Campylobacter pylori J99)

FRDA_HELPY (O06913)

Fumarate reductase flavoprotein subunit (EC 1.3.99.1). {GENE: FRDA OR HP0192} - Helicobacter pylori (Campylobacter pylori)

GCP_HELPJ (Q9ZJ27)

Probable O-sialoglycoprotein endopeptidase (EC 3.4.24.57) (Glycoprotease). {GENE: GCP OR JHP1491} - Helicobacter pylori J99 (Campylobacter pylori J99)

GCP_HELPY (P55996)

Probable O-sialoglycoprotein endopeptidase (EC 3.4.24.57) (Glycoprotease).

{GENE: GCP OR HP1584} - Helicobacter pylori (Campylobacter pylori)

GIDA_HELPJ (Q9ZML9)

Glucose inhibited division protein A. {GENE: GIDA OR JHP0199} -
Helicobacter pylori J99 (Campylobacter pylori J99)

GIDA_HELPY (P56138)

Glucose inhibited division protein A. {GENE: GIDA OR HP0213} -
Helicobacter pylori (Campylobacter pylori)

GPDA_HELPJ (Q9ZKP0)

Glycerol-3-phosphate dehydrogenase [NAD(P)+] (EC 1.1.1.94)
(NAD(P)H-dependent glycerol-3-phosphate dehydrogenase). {GENE: GPSA
OR JHP0895} - Helicobacter pylori J99 (Campylobacter pylori J99)

GPDA_HELPY (O25614)

Glycerol-3-phosphate dehydrogenase [NAD(P)+] (EC 1.1.1.94)
(NAD(P)H-dependent glycerol-3-phosphate dehydrogenase). {GENE: GPSA
OR HP0961} - Helicobacter pylori (Campylobacter pylori)

HCPA_HELPJ (Q9ZMM1)

Beta-lactamase hcpA precursor (EC 3.5.2.6) (Cysteine-rich 28 kDa protein).
{GENE: HCPA OR JHP0197} - Helicobacter pylori J99 (Campylobacter
pylori J99)

HCPA_HELPY (O25001)

Beta-lactamase hcpA precursor (EC 3.5.2.6) (Cysteine-rich 28 kDa protein).
{GENE: HCPA OR HP0211} - Helicobacter pylori (Campylobacter pylori)

HEMN_HELPJ (Q9ZLH0)

Oxygen-independent coproporphyrinogen III oxidase (EC 1.-.-.-)
(Coproporphyrinogenase) (Coprogen oxidase). {GENE: HEMN OR JHP0610} -
Helicobacter pylori J99 (Campylobacter pylori J99)

HEMN_HELPY (O25376)

Oxygen-independent coproporphyrinogen III oxidase (EC 1.-.-.-)
(Coproporphyrinogenase) (Coprogen oxidase). {GENE: HEMN OR HP0665} -
Helicobacter pylori (Campylobacter pylori)

HMCT_HELPJ (Q9ZL53)

Cadmium, zinc and cobalt transporting ATPase (EC 3.6.3.3) (EC 3.6.3.5).
{GENE: CADA OR JHP0727} - Helicobacter pylori J99 (Campylobacter
pylori J99)

HMCT_HELPY (Q59465)

Cadmium, zinc and cobalt transporting ATPase (EC 3.6.3.3) (EC 3.6.3.5).
{GENE: CADA OR HP0791} - Helicobacter pylori (Campylobacter pylori)

KDGL_HELPJ (Q9ZLE0)

Diacylglycerol kinase (EC 2.7.1.107) (DAGK) (Diglyceride kinase) (DGK).
{GENE: DGKA OR JHP0640} - Helicobacter pylori J99 (Campylobacter pylori J99)

KDGL_HELPY (P56411)

Diacylglycerol kinase (EC 2.7.1.107) (DAGK) (Diglyceride kinase) (DGK).
{GENE: DGKA OR HP0700} - Helicobacter pylori (Campylobacter pylori)

LEP_HELPJ (Q9ZLQ5)

Signal peptidase I (EC 3.4.21.89) (SPase I) (Leader peptidase I). {GENE: LEPB OR JHP0523} - Helicobacter pylori J99 (Campylobacter pylori J99)

LEP_HELPY (O25300)

Signal peptidase I (EC 3.4.21.89) (SPase I) (Leader peptidase I). {GENE: LEPB OR HP0576} - Helicobacter pylori (Campylobacter pylori)

LSPA_HELPJ (Q9ZMZ3)

Lipoprotein signal peptidase (EC 3.4.23.36) (Prolipoprotein signal peptidase) (Signal peptidase II) (SPase II). {GENE: LSPA OR JHP0069} - Helicobacter pylori J99 (Campylobacter pylori J99)

LSPA_HELPY (P25178)

Lipoprotein signal peptidase (EC 3.4.23.36) (Prolipoprotein signal peptidase) (Signal peptidase II) (SPase II). {GENE: LSPA OR URED OR HP0074} - Helicobacter pylori (Campylobacter pylori)

MTN_HELPJ (Q9ZMY2)

MTA/SAH nucleosidase [Includes: 5'-methylthioadenosine nucleosidase (EC 3.2.2.16); S-adenosylhomocysteine nucleosidase (EC 3.2.2.9)]. {GENE: MTN OR PFS OR JHP0082} - Helicobacter pylori J99 (Campylobacter pylori J99)

MTN_HELPY (O24915)

MTA/SAH nucleosidase [Includes: 5'-methylthioadenosine nucleosidase (EC 3.2.2.16); S-adenosylhomocysteine nucleosidase (EC 3.2.2.9)]. {GENE: MTN OR PFS OR HP0089} - Helicobacter pylori (Campylobacter pylori)

NADA_HELPJ (Q9ZJN1)

Quinolinate synthetase A. {GENE: NADA OR JHP1274} - Helicobacter pylori J99 (Campylobacter pylori J99)

NADA_HELPY (O25910)

Quinolinate synthetase A. {GENE: NADA OR HP1356} - Helicobacter pylori (Campylobacter pylori)

PYRD_HELPJ (Q9ZM11)

Dihydroorotate dehydrogenase (EC 1.3.3.1) (Dihydroorotate oxidase) (DHODEhase) (DHODase) (DHOD). {GENE: PYRD OR JHP0412} - Helicobacter pylori J99 (Campylobacter pylori J99)

PYRD_HELPY (O25655)

Dihydroorotate dehydrogenase (EC 1.3.3.1) (Dihydroorotate oxidase) (DHODEhase) (DHODase) (DHOD). {GENE: PYRD OR HP1011} - Helicobacter pylori (Campylobacter pylori)

PYRX_HELPJ (Q9ZMG9)

Probable dihydroorotase-like protein (Aspartate carbamoyltransferase 42 kDa non-catalytic chain). {GENE: PYRC' OR JHP0251} - Helicobacter pylori J99 (Campylobacter pylori J99)

PYRX_HELPY (O25045)

Probable dihydroorotase-like protein (Aspartate carbamoyltransferase 42 kDa non-catalytic chain). {GENE: PYRC' OR HP0266} - Helicobacter pylori (Campylobacter pylori)

RADA_HELPJ (Q9ZMK9)

DNA repair protein radA homolog (DNA repair protein sms homolog). {GENE: RADA OR SMS OR JHP0209} - Helicobacter pylori J99 (Campylobacter pylori J99)

RADA_HELPY (P56148)

DNA repair protein radA homolog (DNA repair protein sms homolog). {GENE: RADA OR SMS OR HP0223} - Helicobacter pylori (Campylobacter pylori)

RIR1_HELPJ (Q9ZLF9)

Ribonucleoside-diphosphate reductase alpha chain (EC 1.17.4.1) (Ribonucleotide reductase). {GENE: NRDA OR JHP0621} - Helicobacter pylori J99 (Campylobacter pylori J99)

RIR1_HELPY (P55982)

Ribonucleoside-diphosphate reductase alpha chain (EC 1.17.4.1) (Ribonucleotide reductase). {GENE: NRDA OR HP0680} - Helicobacter pylori (Campylobacter pylori)

RODA_HELPJ (Q9ZLA0)

Rod shape-determining protein rodA. {GENE: MRDB OR RODA OR JHP0680} - Helicobacter pylori J99 (Campylobacter pylori J99)

RODA_HELPY (P56098)

Rod shape-determining protein rodA. {GENE: MRDB OR RODA OR HP0743} - Helicobacter pylori (Campylobacter pylori)

RUVA_HELPJ (Q9ZKW9)

Holliday junction DNA helicase ruvA. {GENE: RUVA OR JHP0815} - Helicobacter pylori J99 (Campylobacter pylori J99)

RUVA_HELPY (O25549)

Holliday junction DNA helicase ruvA. {GENE: RUVA OR HP0883} - Helicobacter pylori (Campylobacter pylori)

RUVB_HELPJ (Q9ZM57)

Holliday junction DNA helicase ruvB. {GENE: RUVB OR JHP0366} - Helicobacter pylori J99 (Campylobacter pylori J99)

RUVB_HELPY (O25699)

Holliday junction DNA helicase ruvB. {GENE: RUVB OR HP1059} - Helicobacter pylori (Campylobacter pylori)

RUVC_HELPJ (Q9ZKX3)

Crossover junction endodeoxyribonuclease ruvC (EC 3.1.22.4) (Holliday junction nuclease ruvC) (Holliday junction resolvase ruvC). {GENE: RUVC OR JHP0811} - Helicobacter pylori J99 (Campylobacter pylori J99)

RUVC_HELPY (O25544)

Crossover junction endodeoxyribonuclease ruvC (EC 3.1.22.4) (Holliday junction nuclease ruvC) (Holliday junction resolvase ruvC). {GENE: RUVC OR HP0877} - Helicobacter pylori (Campylobacter pylori)

RUVX_HELPJ (Q9ZMA5)

Putative Holliday junction resolvase (EC 3.1.-.-). {GENE: JHP0317} - Helicobacter pylori J99 (Campylobacter pylori J99)

RUVX_HELPY (O25101)

Putative Holliday junction resolvase (EC 3.1.-.-). {GENE: HP0334} - Helicobacter pylori (Campylobacter pylori)

SDHL_HELPJ (Q9ZMU7)

L-serine dehydratase (EC 4.3.1.17) (L-serine deaminase) (SDH) (L-SD). {GENE: SDAA OR JHP0120} - Helicobacter pylori J99 (Campylobacter

pylori J99)

SDHL_HELPY (P56072)

L-serine dehydratase (EC 4.3.1.17) (L-serine deaminase) (SDH) (L-SD).

{GENE: SDAA OR HP0132} - Helicobacter pylori (Campylobacter pylori)

TPX_HELPJ (Q9ZKE7)

Probable thiol peroxidase (EC 1.11.1.-). {GENE: TPX OR JHP0991} -

Helicobacter pylori J99 (Campylobacter pylori J99)

TPX_HELPY (O25151)

Probable thiol peroxidase (EC 1.11.1.-). {GENE: TPX OR HP0390} -

Helicobacter pylori (Campylobacter pylori)

TSAA_HELPJ (P56876)

Probable peroxiredoxin (26 kDa antigen). {GENE: TSAA OR JHP1471} -

Helicobacter pylori J99 (Campylobacter pylori J99)

TSAA_HELPY (P21762)

Probable peroxiredoxin (26 kDa antigen). {GENE: TSAA OR HP1563} -

Helicobacter pylori (Campylobacter pylori)

Search in TrEMBL: There are matches to 95 out of 1016356 entries

O24946

Serine transporter (SDAC) {GENE:HP0133} - Helicobacter pylori
(Campylobacter pylori)

O24955

Cytochrome C oxidase, heme B and copper-binding subunit,
membrane-bound (EC 1.9.3.1) (FIXN) {GENE:HP0144} - Helicobacter pylori
(Campylobacter pylori)

O24956

Cytochrome C oxidase, MONOHEME subunit, membrane-bound (FIXO)
{GENE:HP0145} - Helicobacter pylori (Campylobacter pylori)

O24957

CBB3-type cytochrome C oxidase subunit Q (CCOQ) {GENE:HP0146} -
Helicobacter pylori (Campylobacter pylori)

O24958

Cytochrome C oxidase, diheme subunit, membrane-bound (FIXP)
{GENE:HP0147} - Helicobacter pylori (Campylobacter pylori)

O25002

Succinyl-diaminopimelate desuccinylase (DAPE) {GENE:HP0212} -
Helicobacter pylori (Campylobacter pylori)

O25044

Cytochrome C biogenesis protein (CCDA) {GENE:HP0265} - Helicobacter
pylori (Campylobacter pylori)

O25143

Protoporphyrinogen oxidase (HEMK) {GENE:HP0381} - Helicobacter pylori
(Campylobacter pylori)

O25216

Oligoendopeptidase F (PEPF) {GENE:HP0470} - Helicobacter pylori
(Campylobacter pylori)

O25219

Molybdenum ABC transporter, periplasmic molybdate-binding protein
(MODA) {GENE:HP0473} - Helicobacter pylori (Campylobacter pylori)

O25250

Glycolate oxidase subunit (GLCD) {GENE:HP0509} - Helicobacter pylori
(Campylobacter pylori)

O25298

Galactosidase acetyltransferase (LACA) {GENE:HP0574} - Helicobacter
pylori (Campylobacter pylori)

O25343

Tetrahydrodipicolinate N-succinyltransferase (DAPD) {GENE:HP0626} -
Helicobacter pylori (Campylobacter pylori)

O25347

Modulator of drug activity (MDA66) {GENE:HP0630} - Helicobacter pylori
(Campylobacter pylori)

O25348

Quinone-reactive NI/FE hydrogenase, small subunit (HYDA)
{GENE:HP0631} - Helicobacter pylori (Campylobacter pylori)

O25387

Methylated-DNA--protein-cysteine methyltransferase (DAT1)

{GENE:HP0676} - Helicobacter pylori (Campylobacter pylori)

O25399

Acetyl coenzyme A acetyltransferase (Thiolase) (FADA) {GENE:HP0690} - Helicobacter pylori (Campylobacter pylori)

O25402

Hydantoin utilization protein A (HYUA) {GENE:HP0695} - Helicobacter pylori (Campylobacter pylori)

O25403

N-METHYLHYDANTOINASE {GENE:HP0696} - Helicobacter pylori (Campylobacter pylori)

O25464

N-acetylmuramoyl-L-alanine amidase (AMIA) {GENE:HP0772} - Helicobacter pylori (Campylobacter pylori)

O25596

D-alanine glycine permease (DAGA) {GENE:HP0942} - Helicobacter pylori (Campylobacter pylori)

O25729

2-keto-3-deoxy-6-phosphogluconate aldolase (EDA) {GENE:HP1099} - Helicobacter pylori (Campylobacter pylori)

O25743

Gamma-glutamyltranspeptidase (GGT) {GENE:HP1118} - Helicobacter pylori (Campylobacter pylori)

O25824

Oxygen-independent coproporphyrinogen III oxidase (HEMN) {GENE:HP1226} - Helicobacter pylori (Campylobacter pylori)

O25989

60 kDa inner-membrane protein {GENE:HP1450} - Helicobacter pylori (Campylobacter pylori)

O25997

Cytochrome C551 peroxidase {GENE:HP1461} - Helicobacter pylori (Campylobacter pylori)

O26019

Proline dipeptidase (PEPQ) {GENE:HP1485} - Helicobacter pylori (Campylobacter pylori)

O32633

coli cells. Two proteins of molecular mass identical to the B. subtilis IF2 alpha and -beta were over-expressed and purified using a new three-step ion-exchange chromatography procedure. The **N - terminal amino acid** sequence of the two proteins was determined and the results confirmed that the two forms were IF2 alpha and -beta, both encoded by the infB gene. The **N - terminal amino acid** sequence determined for IF2 beta is Met94-Gln-Asn-Asn-Gln-Phe. The presence of methionine at position 94 shows that this form is, in fact, the result of a second translational initiation in infBB.su mRNA, since the codon at amino acid position 94 is GUG, which is the normal codon for valine, but also known to be an initiator codon. This is a new example of the unusual tandem translation in E. coli of an open mRNA reading frame.

Tags: Support, Non-U.S. Gov't

Descriptors: Bacillus subtilis--genetics--GE; *Bacterial Proteins --genetics--GE; *Escherichia coli--genetics--GE; * **Peptide Initiation Factors** --genetics--GE; *Translation, Genetic; Amino Acid Sequence; Bacterial Proteins--metabolism--ME; Base Sequence; Chromatography, Gel; Cloning, Molecular; DNA, Bacterial; Electrophoresis, Polyacrylamide Gel; Molecular Sequence Data; Open Reading Frames; **Peptide Initiation Factors** --isolation and purification--IP; **Peptide Initiation Factors** --metabolism --ME; **Prokaryotic Initiation Factor-2**

CAS Registry No.: 0 (Bacterial Proteins); 0 (DNA, Bacterial); 0 (Peptide Initiation Factors); 0 (Prokaryotic Initiation Factor-2)

Gene Symbol: infB

Record Date Created: 19921207

Record Date Completed: 19921207

7/9/14

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2003 The Dialog Corp. All rts. reserv.

07244546 92107175 PMID: 1729602

The **suil** suppressor locus in *Saccharomyces cerevisiae* encodes a translation factor that functions during tRNA(iMet) recognition of the start codon.

Yoon H J; Donahue T F

Department of Biology, Indiana University, Bloomington 47405.

Molecular and cellular biology (UNITED STATES) Jan 1992, 12 (1) p248-60, ISSN 0270-7306 Journal Code: 8109087

Contract/Grant No.: GM32263; GM; NIGMS

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

We initiated a genetic reversion analysis at the HIS4 locus to identify components of the translation initiation complex that are important for ribosomal recognition of an initiator codon. Three unlinked suppressor loci, **suil**, **sui2**, and **SUI3**, that restore expression of both HIS4 and HIS4-lacZ in the absence of an AUG initiator codon were identified. In previous studies, it was demonstrated that the **sui2** and **SUI3** genes encode mutated forms of the alpha and beta subunits, respectively, of eukaryotic translation initiation factor 2 (eIF-2). In this report, we describe the molecular and biochemical characterizations of the **suil** suppressor locus. The DNA sequence of the **SUI1+** gene shows that it encodes a protein of 108 amino acids with a calculated Mr of 12,300. The **suil** suppressor genes all contain single base pair changes that alter a single amino acid within this 108-amino-acid sequence. **suil** suppressor strains that are temperature sensitive for growth on enriched medium have altered polysome profiles at the restrictive temperature typical of those caused by alteration of a protein that functions during the translation initiation process. Gene disruption experiments showed that the **SUI1+** gene encodes an essential protein, and **antibodies** directed against the **SUI1+** coding region identified a protein with the predicted Mr in a ribosomal salt wash fraction. As observed for **sui2** and **SUI3** suppression events, protein sequence analysis of His4-beta-galactosidase fusion proteins produced by **suil** suppression events indicated that a UUG codon is used as the site of translation initiation in the absence of an AUG start codon in HIS4.

Changing the penultimate proline codon 3' to UUG at his4 to a Phe codon (UUC) blocks aminopeptidase cleavage of the **amino - terminal amino** acid of the His4-beta-galactosidase protein, as noted by the appearance of Met in the first cycle of the Edman degradation reaction. The appearance of Met in the first cycle, as noted, in either a suil or a SUI3 suppressor strain showed that the mechanism of suppression is the same for both suppressor genes and allows the initiator tRNA to mismatch base pair with the UUG codon. This suggests that the Suil gene product performs a function similar to that of the beta subunit of eIF-2 as encoded by the SUI3 gene. However, the Suil gene product does not appear to be a required subunit of eIF-2 on the basis of purification schemes designed to identify the GTP-dependent binding activity of eIF-2 for the initiator tRNA. (ABSTRACT TRUNCATED AT 400 WORDS)

Tags: Support, U.S. Gov't, P.H.S.

Descriptors: Fungal Proteins--genetics--GE; *Genes, Suppressor; * **Peptide Initiation Factors** --genetics--GE; *RNA, Transfer, Met--genetics--GE; *Saccharomyces cerevisiae--genetics--GE; Amino Acid Sequence; Base Sequence; Blotting, Western; Codon; DNA, Fungal; Eukaryotic Initiation Factor-2 --metabolism--ME; Fungal Proteins--metabolism--ME; Genes, Fungal; Molecular Sequence Data; Mutation; **Peptide Initiation Factors** --metabolism--ME; RNA, Fungal; Restriction Mapping; Sequence Alignment; Translation, Genetic Molecular Sequence Databank No.: GENBANK/M77514; GENBANK/X51910

CAS Registry No.: 0 (Codon); 0 (DNA, Fungal); 0 (Eukaryotic Initiation Factor-2); 0 (Fungal Proteins); 0 (Peptide Initiation Factors); 0 (RNA, Fungal); 0 (RNA, Transfer, Met); 144814-03-9 (suil protein)

Record Date Created: 19920210

Record Date Completed: 19920210

7/9/15

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2003 The Dialog Corp. All rts. reserv.

06734952 90360989 PMID: 2390971

Cloning and expression of eukaryotic initiation factor 4B cDNA: sequence determination identifies a common RNA recognition motif.

Milburn S C; Hershey J W; Davies M V; Kelleher K; Kaufman R J

Genetics Institute, Cambridge, MA 02140.

EMBO journal (ENGLAND) Sep 1990, 9 (9) p2783-90, ISSN 0261-4189

Journal Code: 8208664

Contract/Grant No.: GM22135; GM; NIGMS

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

Eukaryotic protein synthesis initiation factor 4B (eIF-4B) is an 80,000 dalton polypeptide which is essential for the binding of mRNA to ribosomes. A highly purified preparation of eIF-4B from HeLa cells was subjected to enzymatic cleavage and **amino - terminal amino** acid sequence analysis. Degenerate oligonucleotide probes were used to isolate a 3851 bp cDNA encoding eIF-4B from a human cDNA library. The DNA encodes a protein comprising 611 residues with a mass of 69,843 daltons. The **amino - terminal** domain of eIF-4B contains a consensus RNA binding domain present in a number of other RNA binding proteins. Expression of eIF-4B in transfected COS-1 cells yielded a polypeptide which reacted with anti-eIF-4B **antiserum** and comigrated with purified eIF-4B. Expression of eIF-4B in COS-1 cells resulted in a general inhibition of translation, possibly due to a 50-fold eIF-4B overproduction.

Tags: Animal; Comparative Study; Human; Support, U.S. Gov't, P.H.S.

Descriptors: **Peptide Initiation Factors** --genetics--GE; Amino Acid Sequence; Base Sequence; Binding Sites; Cell Line; Cloning, Molecular; Gene Expression; Hela Cells--metabolism--ME; Molecular Sequence Data; Oligonucleotide Probes; Peptide Fragments--isolation and purification--IP; Protein Conformation; RNA, Messenger--genetics--GE; RNA, Messenger --metabolism--ME; Sequence Homology, Nucleic Acid; Transfection

CAS Registry No.: 0 (Oligonucleotide Probes); 0 (Peptide Fragments); 0 (Peptide Initiation Factors); 0 (RNA, Messenger); 0 (eIF-4B)

Record Date Created: 19901004
Record Date Completed: 19901004

7/9/16

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2003 The Dialog Corp. All rts. reserv.

06611439 90236932 PMID: 2110148

Isolation and molecular genetic characterization of the Bacillus subtilis gene (infB) encoding protein synthesis initiation factor 2.

Shazand K; Tucker J; Chiang R; Stansmore K; Sperling-Petersen H U; Grunberg-Manago M; Rabinowitz J C; Leighton T

Institut de Biologie Physico-Chimique, Paris, France.

Journal of bacteriology (UNITED STATES) May 1990, 172 (5) p2675-87,
ISSN 0021-9193 Journal Code: 2985120R

Contract/Grant No.: AMO2109; AM; NIADDK

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

Western blot (immunoblot) analysis of Bacillus subtilis cell extracts detected two proteins that cross-reacted with monospecific polyclonal antibody raised against Escherichia coli initiation factor 2 alpha (IF2 alpha). Subsequent Southern blot analysis of B. subtilis genomic DNA identified a 1.3-kilobase (kb) HindIII fragment which cross-hybridized with both E. coli and Bacillus stearothermophilus IF2 gene probes. This DNA was cloned from a size-selected B. subtilis plasmid library. The cloned HindIII fragment, which was shown by DNA sequence analysis to encode the N-terminal half of the B. subtilis IF2 protein and 0.2 kb of upstream flanking sequence, was utilized as a homologous probe to clone an overlapping 2.76-kb ClaI chromosomal fragment containing the entire IF2 structural gene. The HindIII fragment was also used as a probe to obtain overlapping clones from a lambda gt11 library which contained additional upstream and downstream flanking sequences. Sequence comparisons between the B. subtilis IF2 gene and the other bacterial homologs from E. coli, B. stearothermophilus, and Streptococcus faecium displayed extensive nucleic acid and protein sequence homologies. The B. subtilis infB gene encodes two proteins, IF2 alpha (78.6 kilodaltons) and IF2 beta (68.2 kilodaltons); both were expressed in B. subtilis and E. coli. These two proteins cross-reacted with antiserum to E. coli IF2 alpha and were able to complement in vivo an E. coli infB gene disruption. Four-factor recombination analysis positioned the infB gene at 145 degrees on the B. subtilis chromosome, between the polC and spcB loci. This location is distinct from those of the other major ribosomal protein and rRNA gene clusters of B. subtilis.

Tags: Comparative Study; Support, Non-U.S. Gov't; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.

Descriptors: Bacillus subtilis--genetics--GE; *Genes, Structural, Bacterial; * Peptide Initiation Factors --genetics--GE; Amino Acid Sequence; Antibodies; Base Sequence; Blotting, Western; Cloning, Molecular; Coliphages--genetics--GE; Cross Reactions; DNA, Bacterial--genetics--GE; DNA, Bacterial--isolation and purification--IP; Escherichia coli--genetics--GE; Genetic Complementation Test; Genotype; Molecular Sequence Data; Plasmids; Prokaryotic Initiation Factor-2; Restriction Mapping; Sequence Homology, Nucleic Acid

Molecular Sequence Databank No.: GENBANK/M34836

CAS Registry No.: 0 (Antibodies); 0 (DNA, Bacterial); 0 (Peptide Initiation Factors); 0 (Plasmids); 0 (Prokaryotic Initiation Factor-2)

Record Date Created: 19900605

Record Date Completed: 19900605

7/9/17

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2003 The Dialog Corp. All rts. reserv.

06271266 89287323 PMID: 2660906

The two forms of the beta-subunit of initiation factor-2 from reticulocyte lysates arise from proteolytic degradation.

Price N T; Nakielny S F; Clark S J; Proud C G

Department of Biochemistry, School of Medical Sciences, University of Bristol, U.K.

Biochimica et biophysica acta (NETHERLANDS) Jul 7 1989, 1008 (2)

p177-82, ISSN 0006-3002 Journal Code: 0217513

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

Dholakia and Wahba (J. Biol. Chem. (1987) 262, 10164-10170) have reported that preparations of purified initiation factor-2 (eIF-2) from rabbit reticulocytes contain two forms of the beta-subunit. These forms differ in their apparent molecular weights as judged by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE), and are accordingly termed beta H (heavy, the slower-migrating species, apparent Mr = 54,300) and beta L (light, the faster-migrating species, apparent Mr = 53,100). We confirm that two forms of eIF-2 beta are present in such preparations, but present evidence that the beta L is generated from beta H during the isolation procedure. Crude reticulocyte lysates contain only the beta H species as judged from immunoblotting of reticulocyte proteins resolved by SDS-PAGE using an **antisera** against eIF-2 beta. The beta L species appears after the ammonium sulphate fractionation step used early in the purification procedure, but is not apparent if a cocktail of proteinase inhibitors is included in the buffers used during the purification, indicating that it is a proteolytic degradation product generated during the isolation procedure. Cleveland mapping failed to reveal any differences between the two species. Both the beta H and the beta L forms are phosphorylated by casein kinase-2, and, as judged by one- and two-dimensional peptide mapping, at identical sites in each species. Since casein kinase-2 phosphorylates serine-2 in eIF-2 beta, the beta L form must still contain the **N - terminal** region and is presumably produced by limited proteolysis at the carboxyl terminus of the beta-subunit.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: Endopeptidases--metabolism--ME; * **Peptide Initiation Factors** --metabolism--ME; *Proteins--metabolism--ME; Blotting, Western; Chromatography--methods--MT; Electrophoresis, Polyacrylamide Gel; Eukaryotic Initiation Factor-2; Molecular Weight; **Peptide Initiation Factors** --isolation and purification--IP; Peptide Mapping; Phosphorylation; Protease Inhibitors; Protein Kinases--metabolism--ME; Proteins--isolation and purification--IP; Rabbits

CAS Registry No.: 0 (Eukaryotic Initiation Factor-2); 0 (Peptide Initiation Factors); 0 (Protease Inhibitors); 0 (Proteins)

Enzyme No.: EC 2.7.1.- (casein kinase); EC 2.7.1.37 (Protein Kinases); EC 3.4.- (Endopeptidases)

Record Date Created: 19890810

Record Date Completed: 19890810

DapE {GENE:DAPE} - Helicobacter pylori (Campylobacter pylori)

O33991

Nucleosidase homolog (Fragment) - Helicobacter pylori (Campylobacter pylori)

O50170

Putative 32.7 kDa protein - Helicobacter pylori (Campylobacter pylori)

O68226

Flavodoxin A {GENE:FLDA} - Helicobacter pylori (Campylobacter pylori)

O87193

Cb-type cytochrome c oxidase subunit I (EC 1.9.3.1) {GENE:CCON} - Helicobacter pylori (Campylobacter pylori)

O87194

Cb-type cytochrome c oxidase subunit II (EC 1.9.3.1) {GENE:CCOO} - Helicobacter pylori (Campylobacter pylori)

O87195

Cb-type cytochrome c oxidase subunit IV (EC 1.9.3.1) {GENE:CCOQ} - Helicobacter pylori (Campylobacter pylori)

O87196

Cb-type cytochrome c oxidase subunit III (EC 1.9.3.1) {GENE:CCOP} - Helicobacter pylori (Campylobacter pylori)

P94843

Putative glycolate oxidase subunit (EC 1.1.3.15) {GENE:GLCD} - Helicobacter pylori (Campylobacter pylori)

P94847

48 kDa antigen - Helicobacter pylori (Campylobacter pylori)

P94848

29 kDa protein - Helicobacter pylori (Campylobacter pylori)

Q83YF0

NudA (Fragment) {GENE:NUDA} - Helicobacter pylori (Campylobacter pylori)

Q83YF1

NudA (Fragment) {GENE:NUDA} - Helicobacter pylori (Campylobacter pylori)

Q83YF2

NudA (Fragment) {GENE:NUDA} - Helicobacter pylori (Campylobacter pylori)

pylori)

Q83YF3

NudA (Fragment) {GENE:NUDA} - Helicobacter pylori (Campylobacter pylori)

Q84DD1

Gamma-glutamyltranspeptidase - Helicobacter pylori (Campylobacter pylori)

Q8GDA0

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA1

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA2

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA3

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA4

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA5

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA6

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA7

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA8

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter pylori (Campylobacter pylori)

Q8GDA9

Conserved hypothetical protein (Fragment) {GENE:HP519} - Helicobacter

pylori (*Campylobacter pylori*)

Q8RNU2

Heat shock protein B subunit (60 kDa chaperonin) (Protein Cpn60) (groEL protein) {GENE:HSPB} - *Helicobacter pylori* (*Campylobacter pylori*)

Q8RNU3

Heat shock protein A subunit (10 kDa chaperonin) (Protein Cpn10) (groES, protein) {GENE:HSPA} - *Helicobacter pylori* (*Campylobacter pylori*)

Q8RNU7

20 kDa membrane-associated lipoprotein (Membrane-associated lipoprotein Lpp20) {GENE:LPP20} - *Helicobacter pylori* (*Campylobacter pylori*)

Q93SI8

26,000 kDa protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9F5N9

Gamma-glutamyltranspeptidase - *Helicobacter pylori* (*Campylobacter pylori*)

Q9L7Q1

Flavodoxin {GENE:FLDA} - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R2X6

60 kDa major heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R4J2

31 kDa major NONSELECTIVE porin protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R4W4

25 kDa major heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R4W5

30 kDa major heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R4W6

56 kDa major heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R4W7

66 kDa major heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R510

60 kDa heat shock protein/groEL homolog (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9R5L5

HSP62=62 kDa urease-associated heat shock protein (Fragment) - *Helicobacter pylori* (*Campylobacter pylori*)

Q9S0Q6

Lipoprotein signal peptidase A {GENE:LSPA} - *Helicobacter pylori* (*Campylobacter pylori*)

Q9X749

HopV protein precursor (27 kDa outer membrane protein) {GENE:HOPV OR OMP27} - *Helicobacter pylori* (*Campylobacter pylori*)

Q9ZIQ5

Adhesin-thiol peroxidase TagD {GENE:TAGD} - *Helicobacter pylori* (*Campylobacter pylori*)

Q9ZJB7

Putative component of cation transport for CBB3-type oxidase {GENE:FIXI OR JHP1396} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZJF8

Putative cytochrome C peroxidase {GENE:JHP1354} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZK00

Oxygen-independent coproporphyrinogen III oxidase {GENE:HEMN_2 OR JHP1147} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZK95

Gamma-glutamyltranspeptidase {GENE:GGT OR JHP1046} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZKB4

2-keto-3-deoxy-6-phosphogluconate aldolase {GENE:EDA OR JHP1025} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZKD9

Protoporphyrinogen oxidase {GENE:HEMG OR JHP1000} - *Helicobacter pylori* J99 (*Campylobacter pylori* J99)

Q9ZL71

Putative probable N-acetylmuramoyl-L-alanine amidase {GENE:AMIA OR

JHP0709} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLE7

Putative hydantoin utilization {GENE:JHP0633} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLE8

Putative hydantoin utilization {GENE:JHP0632} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLK9

Putative 2,3,4,5-tetrahydropyridine-2-carboxylate N-succinyltransferase {GENE:DAPD OR JHP0570} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLW7

Putative glycolate oxidase {GENE:GLCD OR JHP0459} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLX5

Putative phospholipase A1 {GENE:PLDA OR JHP0451} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLZ8

Molybdate ABC transporter, ATP-binding protein {GENE:MODC OR JHP0427} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZLZ9

Molybdate ABC transporter, permease {GENE:MODB OR JHP0426} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZM00

Molybdate ABC transporter, periplasmic-binding protein {GENE:MODA OR JHP0425} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZM03

Oligopeptidase {GENE:PEPF OR JHP0422} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZM36

Putative proline peptidase {GENE:PEPQ OR JHP0387} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZME9

ATP-dependent zinc METALLOPEPTIDASE {GENE:FTSH_1 OR JHP0271} - Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMH0

Putative cytochrome C-type biogenesis protein {GENE:CCDA OR JHP0250}
- Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMM0

Succinyl-diaminopimelate desuccinylase {GENE:DAPE OR JHP0198} -
Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMT3

Cytochrome oxidase (CBB3-type) {GENE:FIXP OR JHP0135} - Helicobacter
pylori J99 (Campylobacter pylori J99)

Q9ZMT4

Putative cytochrome oxidase (CBB3-type) {GENE:FIXQ OR JHP0134} -
Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMT5

Cytochrome oxidase (CBB3-type) {GENE:FIXO OR JHP0133} -
Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMT6

Cytochrome oxidase (CBB3-type) {GENE:FIXN OR JHP0132} -
Helicobacter pylori J99 (Campylobacter pylori J99)

Q9ZMU6

L-serine transporter, similar to E. COLI NHAB {GENE:SDAC OR JHP0121}
- Helicobacter pylori J99 (Campylobacter pylori J99)

New Search

in Swiss-Prot/TrEMBL by AC, ID,
description, gene name, organism
**Please do NOT use any boolean
operators (and, or, etc.)**


If you would like to retrieve all the Swiss-Prot/TrEMBL entries contained in this list, you can enter a file name. These entries will then be saved to a file under this name in the directory *outgoing* of the ExPASy anonymous ftp server, from where you can download it. (Please note that this temporary file will only

be kept for 1 week.)

File name:

Format: ☒ Swiss-Prot ☐ Fasta

or

 ExpASy Home page		Site Map		Search ExpASy		Contact us		Swiss-Pr t	
Hosted by NCSC US		Mirror sites:		Bolivia	Canada	China	Korea	Switzerland	Taiwan